WARNING KEEP OUT OF REACH OF CHILDREN



Tordon[™] 2G Gold

GRANULAR HERBICIDE

ACTIVE INGREDIENTS:

Contains 20 g/kg picloram and 0.25 g/kg aminopyralid as potassium salts in the form of a pellet.

GROUP 4 HERBICIDE

Controls many difficult-to-kill perennial weeds such as Ragwort, Thistles, Oxeye Daisy and Giant Buttercup as well as Broom, Gorse, Tutsan, Sweet Brier and Blackberry.

READ THE LABEL BEFORE USE

NET CONTENTS: 5 kg, 20 kg

Corteva Agriscience New Zealand Limited Private Bag 2017 New Plymouth 4342 Telephone 0800 803 939 www.corteva.co.nz



INFORMATION WITHIN THIS BOX IS NOT PART OF THE LABEL

WARNING



Skin sensitisation Category 1 Hazardous to soil organisms

QUALIFICATION REQUIREMENTS

Persons mixing, loading, applying, or otherwise handling Tordon™ 2G Gold must meet the qualification requirements set out in the EPA Hazardous Substances (Hazardous Property Controls) Notice 2017.

RECORD KEEPING

Records of use, as described in the EPA Hazardous Substances (Hazardous Property Controls) Notice 2017 and the New Zealand Standard for the Management of Agrichemicals (NZS 8409), must be kept if more than 3 kg of TordonTM 2G Gold is applied or discharged within 24 hours in a place where members of the public may be present or where the substance may enter air or water and leave the place.

PROHIBITED USES

DO NOT use on turf, including lawn and playing fields.

DO NOT apply into, onto or over water.

HEALTH HAZARDS

May cause an allergic skin reaction. Avoid breathing dust or vapours. DO NOT eat, drink or smoke while using this product. Remove protective clothing immediately after handling this product, and wash hands and face before meals and after work.

ENVIRONMENTAL HAZARDS

Hazardous to soil organisms.

SELECTIVE HERBICIDE - VERY TOXIC TO SOME PLANT SPECIES. Certain plants may be damaged or killed by contact with this product. Do not allow movement outside the target area to occur.

HANDLING PRECAUTIONS

EQUIPMENT

Apply using accurately calibrated and maintained equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS 8409).

STORAGE

Store in the original container tightly closed and in a locked, dry, cool, well-ventilated area, well away from foodstuffs.

Storage must be generally in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS 8409). Stores containing 100 kg or more of Tordon™ 2G Gold are subject to signage. See the Health and Safety at Work (Hazardous Substances) Regulations

INFORMATION WITHIN THIS BOX IS NOT PART OF THE LABEL

and the New Zealand Standard for the Management of Agrichemicals (NZS 8409) for further information.

PERSONAL PROTECTION

When opening the container, preparing the product for use and applying the product, wear chemical resistant gloves, and cotton overalls buttoned to the neck and wrist. Contaminated work clothing should not be allowed outside of the workplace. Take off contaminated clothing and wash it before re-use.

SPILLAGE

DO NOT touch or walk through spilled material. Wear the protective equipment recommended above. Stop further spill occurring and prevent entry into waterways and drains. **Small spills:** In case of spill on floor or paved surfaces sweep up and place in drums for disposal at a licensed local authority landfill. Spill residues may be cleaned up using water and detergent. Absorb and collect washings and place in the same sealable container for disposal. Seek advice from the SDS, product label or Corteva Agriscience regarding disposal. **Large spills:** Report large spills to Corteva Agriscience Emergency Services at 0800 844 455.

DISPOSAL

Triple rinse empty containers. Triple rinsed containers containing the Agrecovery logo on the label, and that are free of all residues (liquid and dry) and have an intact legible label should be taken to an Agrecovery container collection site. Otherwise crush and bury in a suitably approved landfill. DO NOT burn empty containers.

Dispose of the product only by using according to the label, via Agrecovery (visit www.agrecovery.co.nz or call 0800 AGRECOVERY (247 326) for information about the type and quantity of chemical acceptable for disposal), or at an approved landfill. DO NOT burn.

DO NOT dispose of this product or equipment cleaning water down drains or sewers.

FIRST AID

If on skin: Wash with plenty of water. If skin irritation or rash occurs: Get medical advice. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If medical advice is needed, have the product container or label at hand.

TRANSPORT

DO NOT carry more than 3 kg on a passenger service vehicle. Quantities of 3 kg or less carried on a passenger service vehicle must be contained within sealed packaging.

SAFETY DATA SHEET

A Safety Data Sheet for Tordon™ 2G Gold is available from Corteva Agriscience on request. Call Customer Service Toll Free on 0800 803 939 or visit www.corteva.co.nz

NOTICE

Important: You **must** read the warning and conditions given on this label. If they are not acceptable, or if you are in doubt as to the proper handling of this product (including its transport, storage, use, disposal, and requirements for approved handlers [if applicable]) then return the unopened product to the supplier at once.

Warning: This product is subject to controls and could be hazardous if handled incorrectly. The **user is responsible** for ensuring that the handling of this product is in accordance with the law (including the *Hazardous Substances and New Organisms Act 1996* and the *Agricultural Compounds and Veterinary Medicines Act 1997*) and with the directions provided. The Registrant declines all liability (whether in contract, tort (including but not limited to negligence), or otherwise) to the maximum extent permitted by law in relation to the handling of this product in a manner which is contrary to the directions given or to any applicable Acts, Regulations or Guidelines.

Further information regarding the applicable Acts, Regulations and Guidelines for the proper handling of this product may be obtained by contacting EPA New Zealand (www.epa.govt.nz) or the Ministry for Primary Industries (www.foodsafety.govt.nz).

This product is warranted to contain the specified active ingredients within accepted analytical tolerance when packed.

Registered to:

Corteva Agriscience New Zealand Limited

Private Bag 2017 New Plymouth, 4342 Telephone 0800 803 939 www.corteva.co.nz

EMERGENCY RESPONSE

(All hours)
Ring from anywhere in New
Zealand
0800 844 455
or Globally
+64-6-751 2407

IN A TRANSPORT EMERGENCY DIAL 111 FOR POLICE OR FIRE

Registered pursuant to the ACVM Act 1997, No P9023. See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to HSNO Act 1996, EPA Approval Code HSR100945 See www.epa.govt.nz for approval controls.

Date of Manufacture: Batch No. Barcode

APPLICATION METHODS

Applications can be made with granule applicators such as the Weedstick granule applicator or Prill Granule applicator, or alternatively by hand. Always wear waterproof gloves when applying this product by hand. Avoid allowing granules to fall inside the gloves.

SPOT APPLICATION ONLY

TordonTM 2G Gold should be applied evenly over the top of the plant and in the case of brush weeds extending out to 30 cm beyond the dripline. When treating bushes on steep slopes more granules should be applied on the uphill side as some downward leaching will occur.

IMPORTANT: This product must be applied in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS 8409).

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residues Levels for Agricultural Compounds.

Except for animal safety reasons no withholding period is required when used as directed.

ANIMAL SAFETY

Poisonous plants (including but not limited to boxthorn, hemlock, nightshades, ragwort and tutu) may become more palatable after application. If concerned that stock health may be affected, stock should be kept away from the treated area until the plants have died down.

PROTECTION OF THE ENVIRONMENT

DO NOT use on turf or in the home garden.

TordonTM 2G Gold is injurious to legumes and other broadleaf plants. Residues in the soil may prevent their early re-establishment after treatment. Susceptible crops and plants include but are not limited to, flowers, fruit trees, legumes, ornamentals, potatoes, shade trees, squash, tomatoes, vegetables, or vine crops. Susceptible crops and plants may be damaged by this herbicide during both growing and dormant periods. If in doubt regarding other susceptible plants, or for further information about planting susceptible plants into a treated area within 2 years of application consult a Corteva Agriscience representative.

DO NOT allow the granules to contact susceptible plants, including the root zone.

DO NOT apply close to, or on, areas where treated soil or surface water may carry the granules off to areas containing, or are to be planted with, susceptible plants.

DO NOT use plant material that has been treated with Tordon[™] 2G Gold for making compost, mushroom substrate or mulching susceptible crops.

RESISTANCE MANAGEMENT



Tordon™ 2G Gold is a pyridine carboxylate herbicide (Group 4) which has an auxin mimic mode of action. Some naturally occurring individual weeds may become resistant when excessive use of the same herbicide occurs. To maximize the effectiveness of herbicides, products containing alternate modes of action should be used in the spray program. Consult http://resistance.nzpps.org/ for alternate modes of action and the details of resistance management strategies (if any) for the crops listed on the label.

DIRECTIONS FOR USE

Tordon™ 2G Gold is recommended as a spot application for control of scattered brush weeds and broadleaf weeds in areas where spray treatment is impractical or undesirable.

DO NOT USE ON PEAT OR HEAVY CLAY SOILS.

TIME OF APPLICATION

Best results are achieved if TordonTM 2G Gold is applied when favourable growth conditions exist and particularly when sufficient moisture is present to move the TordonTM 2G Gold from the granule to the soil. Applications outside these times may reduce efficacy.

WEEDS	RATE	COMMENTS
Ragwort, nodding thistle, docks.	2 g per plant	Apply half a level teaspoon to the crushed centre of each plant.
Broom, fennel, giant buttercup, goats rue, hemlock, inkweed, lupin, Mexican devil weed, oxeye daisy, St John's wort, woolly nightshade.	30 g/sq.m	Best results are achieved on light sandy or stony soils and when applied late winter to late spring.
Gorse, blackberry, boxthorn (seedling), barberry (small bushes), heath, horehound, Himalayan honeysuckle, sweet briar, matagouri, tutu, tutsan	55 g/sq.m	

[™] Trademark of the Corteva Agriscience and its affiliated companies. © 2023 Corteva.